

09.01.03



REC'D 03 FEB 2003

WIPB OCT

# Kongeriget Danmark

Patent application No.: PA 2002 00031

Date of filing: 09. January 2002

Applicant: Arexis AB  
Anne Lunden

This is to certify the correctness of the following information:

The attached document is a true copy of the following document:

- The specification, claims and drawings as filed with the application on the filing date indicated above.



**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Patent- og Varemærkestyrelsen  
Økonomi- og Erhvervsministeriet

21. January 2003

Bo Z. Tidemann

181

Modtaget PVS  
9 JAN. 2002

1. Gebyrer:

Ansigingsgebyr  
Kravgebyr  
ITS-nyhedsundersøgelises-gebyr

12. Bilagsfortegnelse:

☒ fremmedsproget beskrivelse  
☐ dansk beskrivelse i 1 ekspl.  
☐ sammendrag i 1 ekspl.  
☒ tegninger i 1 ekspl.  
☐ prioritetsdokument  
☐ fuldmagt  
☐ overdragelsesdokument

Fig nr. ønskes  
publiceret sammen med  
sammendraget

15. Behandling af  
fremmedspoget ansøgning  
min. ønskes

norsk  
svensk  
engelsk

# Patentansøgning

Læs venligst vejledningen til de enkelte punkter

2. Fuldmægtigs referencenr.:  
31661 DK 01

3. International indleveringsdag:

International ansøgningensnr:

Kapitel 1

Kapitel II

4. Ansøger (fulde navn og adresse):

Arexis AB  
Krokslätts Fabriker 30  
S-431 37 Mölndal  
Sverige

Flere ansøgere  
på side 2

5. Fuldmægtig (navn og adresse):

Plougmann & Vingtoft a/s  
Sankt Annæ Plads 11  
1250 København K

Telefon: 33 63 93 00

Telefax: 33 63 96 00

6. Opfinder (fornavn, efternavn, adresse)

Anne Lundén  
Funbogården  
S-755 97 Uppsala  
Sverige

6a. Opfinder (fornavn, efternavn, adresse):

☒

Flere  
opfindere på  
side 2

Leif Andersson  
Bergagatan 30  
S-752 39 Uppsala  
Sverige

7. Opfindelsens benævnelse:

Detection of mutations

8. Prioritetspåstand(e):

☐

Flere priori-  
tetspåstande  
på side 2

Dato  
Dato  
Dato

Land  
Land  
Land

Nr.  
Nr.  
Nr.

9. Ansøgningen omfatter deponering af mikroorganisme-kultur(er), som angivet i patentlovens §8a, stk. 1.

10. ☒ Ansøgningen omfatter en sekvensliste.

11. Ansøgningen er fremkommet ved deling eller udskillelse.  
Stamansøgningens nr.: Ansøgt løbedag:

13. Ansøgningen er tidligere indleveret pr. telefax den:

14. Dato og underskrift:  
København, den 9. januar 2002

*Plougmann & Vingtoft A/S*  
Plougmann & Vingtoft A/S

PA 2002 00031

- 9 JAN. 2002

**Detection of mutations****FIELD OF INVENTION**

- 5 The invention relates to mutations in the flavin-containing monooxygenase (FMO3) gene and methods for detection of such mutations in farm animals and poultry.

**BACKGROUND OF INVENTION**

## 10 Occurrence of fishy off-flavor in cow's milk

- Fishy off-flavor in milk is a quality defect recently observed in bulk milk in Sweden, which causes considerable losses to both milk producers and dairy companies. Statistical data from an investigation made 1999 regarding the occurrence of off-flavor in milk showed that out of roughly 2,100 herds, 115 (i.e. > 5%) had received one or more complaints about fishy off-flavor in bulk milk. Altogether, these 115 herds had
- 15 received 242 complaints about a moderate fishy off-flavor, whereas 18 bulk milk samples were classified as having a pronounced fishy odor/taste. Exact figures on the current prevalence of fishy odor among Swedish dairy cows are difficult to estimate as no routine organoleptic testing on milk from individual cows is performed. However, when random samples of bulk milk from individual farms were judged to have a fishy off-flavor, tests on samples from individual cows were sometimes performed. These tests have shown that
- 20 the off-flavor most often originates from one or a few cows within a herd. In order to exceed the threshold above which the test panel perceives a fishy off-flavor in the farm bulk milk, a sufficiently high proportion of the total milk must come from affected cows. Consequently, an individual cow with a fishy off-flavor in the milk will go undetected unless she belongs to a fairly small herd. The above figure of 5% is thus likely an underestimate of the problem. To what extent this off-flavor occurs in other countries is not known, as the
- 25 vast majority do not perform regular tests for off-flavor in bulk milk at the farm level. There are so far no indications that the defect has any fatal effects on traits subjected to natural selection, like calf survival, or traits included in the breeding goal for milk production.

**Characteristics**

- 30 Fishy off-flavor is characterized by a distinct, unpleasant taste and smell, reminiscent of rotting fish. A fishy smell in milk (Humphriss, 1953; Corfield, 1955) and coffee cream (Eyer *et al.* 1990) has previously been reported in connection with bacterial degradation of lecithin, choline, and betaine, the latter two being intermediate products in the breakdown of lecithin to trimethylamine (TMA) oxide with TMA as an intermediate compound. A further potential source of fishy taint was related to selective oxidation of
- 35 butterfat (Swoboda & Peers, 1977). The human nose is extremely sensitive to the TMA odor (Ayesh & Smith, 1990), e.g. the olfactory threshold for detection in milk lies around 1-2 ppm (Metha *et al.* 1974; von Gunten *et al.* 1976). As a consequence, milk from a few affected cows in a herd has been shown to be sufficient to cause a fishy off-flavor to the whole bulk milk. There have been a few reports on fishy odor in milk related to TMA content in milk from cows on wheat pasture (Metha *et al.* 1974; von Gunten *et al.*
- 40 1976; Kim *et al.* 1980).

***Similar phenomena in other species***

Problems with fishy odor associated with elevated TMA levels have been described in human (Fish-odor

syndrome' or 'Trimethylaminuria', OMIM #602079, <http://www.ncbi.nlm.nih.gov>) and chicken (Hobson-Frohock *et al.* 1973). In human, abnormal secretion of TMA has been observed in breath, urine, sweat, saliva, and vaginal secretions (Humbert *et al.* 1970), whereas in chicken the TMA has predominantly been found in egg yolk (Hobson-Frohock *et al.* 1973). The TMA is derived from the intestinal degradation of food/feed components rich in TMA or its precursors. Under normal conditions the TMA produced is oxidized to the odor- and tasteless compound TMA-oxide by the liver enzyme flavin-containing monooxygenase (FMO) (Hlavica & Kehl, 1977). The TMA oxide is thereafter excreted in the urine (Al-Walsh *et al.* 1987a). The fishy odor is a consequence of impaired oxidation of TMA (Pearson *et al.* 1979; Spellacy *et al.* 1979).

#### 10 Genetic causes

The fishy odor shows a recessive mode of inheritance in human (Al-Walsh *et al.* 1987b, 1987c; Ayesh *et al.* 1993) whereas in chicken it is described as 'semidominant' because the expression has been shown to be dependent on the ingestion of e.g. rapeseed meal (Bolton *et al.* 1976). In human, parents heterozygous for the defect were shown to excrete elevated amounts of TMA in the urine when given oral doses of 600 mg TMA (Al-Walsh *et al.* 1987c, 1989; Ayesh *et al.* 1990, 1993). However, in none of the cases did the treatment result in a fishy odor.

The fishy odor syndrome in humans has recently been shown to be due to mutations in *FMO3* encoding an isoform of flavin-containing monooxygenase (Dolphin *et al.* 1997b; Treacy *et al.* 1998; Akerman *et al.* 1999; Basarab *et al.* 1999; Forrest *et al.* 2001). The gene has been localized to chromosome 1q23-q25 (<http://www.ncbi.nlm.nih.gov/LocusLink>, August 2001) and its genomic sequence is known (Dolphin *et al.* 1997a). The human *FMO3* gene contains one non-coding and 8 coding exons and is 22.5 kb long. (GenBank AH006707, GenBank AL021026)

#### Dietary causes

The elevated concentration of TMA observed in various tissues in human, chicken, and cattle is likely to be due to a combination of genetic and dietary effects. Feed components rich in TMA precursors such as betaine, choline and sinapin, are beet products, green leaves and cereals, and rapeseed products. A high intake of these products may result in an accumulation of TMA, which in turn overloads the enzyme system. Rapeseed also contains progoitrin, a substance that in chicken has been shown to act as an FMO inhibitor by competing with TMA for the enzyme's active binding site (Pearson *et al.* 1982). In humans, the major source of TMA is marine fish and other seafoods (Zhang *et al.* 1999).

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. (a) Alignment of codon 236-238 for the normal (R238) and mutant (R238X) *FMO3* allele in cattle. This region was analyzed using pyrosequencing. The location of the sequencing primer and the six positions sequenced are indicated. (b) Pyrosequencing results from the three different genotypes at codon 238. The sequenced reverse strand has the sequence (A/G)AGTGA where the A/G polymorphism corresponds to the R238X nonsense mutation.

#### 40 DESCRIPTION OF THE INVENTION

According to the invention it is shown that the *FMO3* nonsense mutation R238X causes fishy off-flavor in milk. This is the first identified gene that has a profound influence on perceived quality of raw milk. The genotyping method described in this study can now be used by the breeding organisations to eliminate the problem in those breeds where a *FMO3* nonsense mutation is present.

The present invention provides polypeptides which are flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide comprising at least a sequence having at least 85% identity, preferably at least 90% identity, more preferably 95% identity, with the polypeptide sequence SEQ ID NO: 15.

5

The present invention further provides polypeptides which are flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide having at least 85% identity, preferably at least 90% identity, more preferably 95% identity, with the polypeptide sequence SEQ ID NO: 15.

- 10 An examples of such a polypeptide is the bovine FMO3 as shown in SEQ ID NO:15. Further examples of polypeptides of the invention are the sheep and goat FMO3.

The present invention also provides polypeptides which are functionally altered mutants of flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide comprising at least a sequence having at

- 15 least 85% identity, preferably at least 90% identity, more preferably 95% identity, with the polypeptide sequence SEQ ID NO: 15.

An example of such functionally altered mutants is a polypeptide resulting from a deletion of a part of the FMO3 polypeptide, such as the R238X variant of the bovine FMO3.

20

Other examples of polypeptides which are functionally altered mutants of flavin-containing monooxygenase 3 (FMO3) according to the invention, may arise from insertions, deletions, and mis-sense mutations in the FMO3 gene.

- 25 The present invention further provides isolated nucleic acid sequences encoding flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide comprising at least a sequence having at least 85% identity, preferably at least 90% identity, more preferably 95% identity, with the polypeptide sequence SEQ ID NO: 15.

- 30 Examples of such isolated nucleic acids are the cDNA sequence encoding the bovine FMO3 shown in SEQ ID NO: 14, the bovine FMO3 gene, or fragments thereof. Examples of such fragments are exons 3, 6, 7 and 9 shown as SEQ ID Nos: 9, 10, 11, and 12, respectively.

- 35 Nucleic acid sequences of the invention include variants of the FMO3 gene, such as the R238X and E287G variants of the bovine FMO3 gene.

- To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = # of identical positions/total # of positions (e.g., overlapping positions) x 100). In one embodiment the two sequences are the same length.
- 45

- The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) J. Mol. Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a protein molecule of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (1997) Nucleic Acids Res. 25:3389. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules. When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.
- The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, only exact matches are counted.

- The invention also includes specific fragments of a nucleic acid sequence of the invention. "Specific fragments" refers to nucleic acids fragments having a sequence that is found only in the nucleic acid sequences of the invention, and is not found in nucleic acid sequences encoding related polypeptides such as FMO1, FMO2, FMO4, and FMO5.

- "Specifically hybridising fragments" refers to nucleic acid fragments which can hybridize, under stringent conditions only with nucleic acids of the invention, without hybridizing with nucleic acid sequences encoding related polypeptides such as FMO1, FMO2, FMO4, and FMO5.

Said specific fragments or specifically hybridising fragments may for example be used as primers or probes for detecting and/or amplifying nucleic acid sequences encoding FMO3 polypeptides.

- Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Molecular Cloning: A Laboratory Manual, 3rd ed., Sambrook et al. eds., Cold Spring Harbor Laboratory Press, 2001; DNA Cloning: A practical Approach, Glover & Hames eds., Oxford University Press, 1996; Nucleic Acid Hybridization: Essential techniques, Ross ed. Wiley, 1998.

- The invention includes sets of primers comprising at least one primer consisting of a specific fragment or specifically hybridising fragment as defined above.

As shown herein, mutations in the bovine *FMO3* gene may lead to fish off-flavour of milk, caused by an altered metabolism of trimethylamine.

- It is further postulated that mutations in the chicken *FMO3* gene may be associated with fish off-flavour of eggs produced by hens carrying such mutations.

- The invention also includes a method for detecting a mutation in the *FMO3* gene of an animal where the mutation will cause an alteration in the metabolism of trimethylamine leading to a fish off-flavour in a food product of the animal or its offspring, the food product can be e.g., milk or eggs. The method comprises:

- obtaining a nucleic acid sample from the animal;
- determining the presence in said nucleic acid sample of a nucleic acid sequence encoding a mutated *FMO3*.

5 In yet another embodiment the invention provides a method for detecting a nucleic acid sequence comprising a mutation in the *FMO3* gene of an animal where the mutation will cause an alteration in the metabolism of trimethylamine leading to a fish off-flavour in a food product of the animal or its offspring.

The method comprises:

- obtaining a nucleic acid sample from the animal;
- 10 - contacting said nucleic acid sample with a nucleic acid probe spanning said mutation under conditions of specific hybridisation between said probe and the mutant sequence to be detected
- detecting the hybridisation complex.

15 Preferably, the process of the invention further comprises, prior to hybridization, PCR amplification from the nucleic acid sample, of a sequence comprising at least the portion of the *FMO3* sequence wherein the mutation is to be detected.

The food product can be e.g. milk or eggs. The animal can be e.g. a cow, a bull, a sheep, a goat, a hen or a cock.

20

Methods allowing the specific hybridisation of a probe only with a perfectly matching complementary sequence, and useful for the detection of punctual mutations are known in the art. They include for instance Allele Specific PCR (Gibbs, Nucleic Acid Res. 17: 2427-2448, 1989), Allele Specific Oligonucleotide Screening (Saiki et al. Nature 324:163-166, 1989).

25

A mutation in the *FMO3* gene may also be detected through detection of polymorphic markers closely linked to said mutation. The invention also provides means for identifying said polymorphic markers, and more specifically polymorphic markers comprised within a genomic DNA sequence comprising at least a portion of a *FMO3* gene, and up to 500 kb, preferably 300 kb, more preferably up to 100 kb of a 3' and/or of a 5'

30 adjacent sequence.

Said polymorphic markers may be obtained for instance, by screening a genomic DNA library from an animal with a probe specific for the *FMO3* gene, in order to select clones comprising said nucleic acid sequence and flanking genomic sequences, and identifying a polymorphic marker in said flanking genomic sequences. The allele(s) of a polymorphic marker associated with a given mutant allele of the *FMO3* gene may also easily be identified by use of a genomic DNA library from an individual wherein the presence of said mutant allele has previously been detected by hybridisation with a nucleic acid probe of the invention.

35

Polymorphic markers include for instance, single nucleotide polymorphisms (SNP), microsatellites, insertion/deletion polymorphisms and restriction fragment length polymorphisms (RFLP). These polymorphic markers may be identified by comparison of sequences flanking the *FMO3* gene obtained from several individuals. Microsatellites may also be identified by hybridisation with a nucleic acid probe specific of known microsatellite motifs. Once a polymorphic marker has been identified, a DNA segment spanning the polymorphic locus may be sequenced and a set of primers allowing amplification of said DNA segment may be designed. The invention also encompasses said DNA primers.

45

Detection of a mutation in *FMO3* gene may be performed by obtaining a sample of genomic DNA from an animal, amplifying a segment of said DNA spanning a polymorphic marker by PCR using a set of primers of the invention, and detecting in said amplified DNA the presence of an allele of said polymorphic marker associated with said mutation.

5

The invention also provides kits for the practice of the methods of the invention. The kits comprise at least one specific fragment of a nucleic acid of the invention, or at least one nucleic acid fragment able to hybridise with a nucleic acid sequence of the invention. Said nucleic acid may be labelled. The kits may also comprise a set of primers of the invention. They may be used in conjunction with commercially available amplification kits. They may also include positive and/or negative control reactions or markers, molecular weight size markers for gel electrophoresis, and the like.

10



## EXAMPLES

### Example 1. Sequencing of the bovine *FMO3* gene

#### 5 Cattle samples and DNA preparation

The present study included two different groups of dairy cattle. The first material comprised 21 cows of the Swedish Red and White breed (SRB) with information from sensory analyses on the milk. Ten of the cows belong to either of the two experimental herds, Jälla and Kungsängen, at Swedish University of Agricultural Sciences in Uppsala. The remaining cows belong to either of four commercial herds, all of which had received complaints about fishy off-flavor in the bulk milk. In three of the six herds, 'control' cows were chosen that were comparable to the affected cows as regards breed, lactation number, and stage of lactation but were producing milk with a normal taste and smell.

A population study was performed in which bulls and cows from the four dairy breeds in Sweden were genotyped for the observed *FMO3* nonsense mutation. From each of the two major breeds, SRB and Swedish Holstein, 100 individuals were chosen, whereas from the two small breeds, Swedish Polled and Swedish Jersey, 25 and 23 individuals were sampled, respectively. Genomic DNA was prepared from blood samples according to standard protocols.

#### PCR amplification and sequencing of the bovine *FMO3* gene from genomic DNA samples

Exonic parts of the *FMO3* gene were amplified from genomic DNA samples using primers corresponding to sequences well conserved between the human, rabbit, mouse, and rat *FMO3* sequences available in GenBank (Table 1). Two genomic DNA samples were used, one from a cow that had been shown to repeatedly produce milk with a fishy flavor and the other from a control cow producing normal milk.

Table 1. Primer sequences used for PCR amplification of the bovine *FMO3* gene from genomic

DNA.

<i>FMO3</i> region amplified <sup>1</sup>	Forward (F) and reverse (R) primer sequences 5'-3'	
<i>exon 3</i>	F: GACCATGCAGAAGAGGGCAG	SEQ ID NO: 1
	R: CTAAACTGTATGTATTTTCAGGAGGTT	SEQ ID NO: 2
<i>exon 6</i>	F: CATCATCAGCTCCAGAAGTGG	SEQ ID NO: 3
	R: TAAAGGCATCAAGCCATAGTT	SEQ ID NO: 4
<i>exon 7</i>	F: CAGAATCCTGAGGAAAGAGCC	SEQ ID NO: 5
	R: ATTACTTGTGCTGCCACGCG	SEQ ID NO: 6
<i>exon 8-9</i>	F: GATGAATGATATTAATGAGAAAATGG	SEQ ID NO: 7
	R: CCGGTCCCACTGGGTCAG	SEQ ID NO: 8

<sup>1</sup>The numbering of exons is based on the described exon-intron organization for the human *FMO3* gene (Dolphin *et al.* 1997a).

The PCR was performed in 10 µl reactions including 1 x PCR buffer, 1.5 mM MgCl<sub>2</sub>, 0.2 mM of each dNTP, 2.5 pmol of each primer, 25 ng genomic DNA, and 0.35 U AmpliTaq DNA polymerase (Perkin Elmer, Branchburg, NJ, USA). Thermocycling was carried out in a PTC200 machine (MJ Research, Watertown, MA, USA) and included 40 cycles with annealing at 53°C for 30 s and extension at 72°C for 2 min. The denaturation step was 95°C for 1-2 min in the first two cycles, and 94°C for 30 s in the remaining cycles. The products were analyzed by agarose gel electrophoresis (4% Nusieve/Seakern 3:1, FMC Bioproducts, Rockland, ME, USA). The PCR products were directly sequenced with BigDye terminators and an ABI377 Instrument (Perkin-Elmer, Foster City, CA, USA). Sequence analysis and comparison of sequences from the cows with fishy flavored and normal milk, respectively, was performed using the Sequencher software (GENE CODES, Ann Arbor, MI, USA).

#### **Bovine *FMO3* gene sequences**

Bovine *FMO3* sequences corresponding to human *FMO3* exon 3, 6, 7, 9, and intron 8 were generated and are given in SEQ ID Nos: 9-13. Subsequently, the complete coding region of the bovine *FMO3* gene was sequenced from RT-PCR products from liver mRNA. The bovine *FMO3* cDNA sequence is given in SEQ ID NO: 14 and the corresponding polypeptide sequence of bovine *FMO3* is given in SEQ ID NO: 15. The exonic sequences generated from genomic DNA showed 100% identity to the cDNA sequence obtained from the RT-PCR products and *FMO3* specificity was strongly supported by the high sequence similarity to other mammalian *FMO3* sequences. For example, a Blast comparison between the complete coding cattle *FMO3* sequence obtained here and the corresponding human sequence (GenBank NM\_006894) showed 85% and 82% identity at the nucleotide and amino acid levels, respectively.

#### ***Identification of a nonsense mutation in the bovine *FMO3* gene***

Partial cattle *FMO3* sequences obtained from a cow producing milk with strong fishy off-flavor and a cow producing normal milk were compared. The compared sequences included altogether 1522 bp and 808 bp of these represent coding sequence. The coding part includes 268 codons, corresponding to approximately 50% of the protein. The only sequence difference found between the two cows was a C/T nucleotide substitution located at position 62 in exon 6 (SEQ ID NO: 10). The C→T substitution changes the codon for arginine (R) at position 238 to a stop codon (X); the numbering of amino acid positions are based on the human amino acid sequence given in Genbank NM\_006894. This nonsense mutation is thus denoted R238X. The pyrosequencing test confirmed that the affected cow was homozygous for the R238X substitution whereas the control cow was homozygous R/R at residue 238 (Fig. 1).

#### **Example 2. Association between mutations in *FMO3* and the presence of fishy off-flavour in cow's milk.**

##### **Genotyping of the *FMO3* R238X nonsense mutation**

The identified single nucleotide polymorphism (SNP) in exon 6 was genotyped by pyrosequencing (Ronaghi et al. 1998) using a Luc96 Instrument (Pyrosequencing AB, Uppsala, Sweden). This included PCR amplification of a 147 bp product from genomic DNA samples using the forward primer 5'-biotin- GAT GAA GGC TAT CCA TGG GAC (SEQ ID NO: 16) and the reverse primer 5'-TAA AGG CAT CAA GCC GTA GTT CTC (SEQ ID NO: 17). The PCR was performed in 20 µl reactions with concentrations of reagents as above. Thermocycling was carried out in a PE9600 Instrument (Perkin-Elmer, Foster City, CA, USA) and included an

Initial denaturation step at 94°C for 2 min followed by 5 cycles with 94°C for 30 s, 55°C for 30 s, 72°C for 45 s, 43 cycles with 94°C for 20 s, 50°C for 30 s, 72°C for 45 s, and a final extension step at 72°C for 5 min. Pyrosequencing on the forward strand was done using the reverse sequencing primer 5' TGA GGA ATG TTT CAA ATC (SEQ ID NO: 18) and conditions according to the manufacturer's recommendations.

5

***RT-PCR amplification and sequencing of the bovine FMO3 cDNA***

Ten livers from SRB cattle were collected and the pyrosequencing test showed that two were heterozygous carriers of the R238X mutation. One of these livers as well as one liver from a homozygous wild-type cow were used for mRNA preparation, reverse transcription (RT)-PCR amplification and direct sequencing of the coding FMO3 region. Two additional primers were used for this purpose, designed from sequences in the untranslated regions which showed conservation between other mammalian FMO3 sequences. Thus, the 5'UTR primer 5' GGA CTT AGA CAC ACA GAA GAA AAG AAG (SEQ ID NO: 19) and the 3'UTR primer 5' GAG GTG TGA AAT TCT TAT TTT TTA AAT AG (SEQ ID NO: 20) were used in pairs with the reverse and forward pyrosequencing PCR primers, respectively. Thus, the coding sequence was amplified in two overlapping pieces, each including the site of the R238X mutation. Poly A mRNA was isolated from cattle livers using the Quickprep mRNA Micro purification kit (Amersham Pharmacia Biotech, Uppsala, Sweden) and RT-PCR was carried out using the First strand cDNA synthesis kit (Amersham Pharmacia Biotech) with ~200 ng mRNA primed with 0.1 µg random hexamers in a 15 µl reaction volume. Two µl of the completed first strand reaction was used in each PCR reaction with a total volume of 12 µl so that the final concentrations of dNTP, Mg<sup>2+</sup>, primers, and AmpliTaq polymerase were the same as above. The thermocycling conditions were as described above for the sequenced genomic DNA amplicons but the 5' part of the cDNA required a repeated PCR to yield enough product for sequencing.

The R238X substitution is closely associated with the presence of fishy off-flavor in cow's milk

25 The pyrosequencing test was used to test a case-control material comprising animals producing milk classified as having a fishy off-flavor and control animals from the same herds (Table 2). A Class 2 score means that two trained persons both classified the sample as having a strong fishy off-flavor whereas Class 1 implies that the two persons judged the fishy off-flavor as strong/moderate, moderate/moderate or moderate/normal. The results revealed a very strong association between the FMO3 R238X nonsense mutation and strong fishy-off flavor since eight out of nine animals with the Class 2 score were homozygous for this mutation and the ninth was heterozygous. In contrast, the five control samples were homozygous normal. The Class 1 group contained animals of all three genotype classes.

Table 2. *FM03* genotype distributions in a group of 21 Swedish Red and White cows with or without fishy off-flavor in milk and in animals representing the four different dairy breeds in Sweden

5	FM03 genotype at codon 238				
	R/R	R/X	X/X	Total	
<b><u>Case/control study</u></b>					
10	<b><i>Sensory evaluation:<sup>1</sup></i></b>				
	Normal	5	0	0	5
	Class 1	3	3	1	7
	Class 2	0	1	8	9
15	<b><u>Population study</u></b>				
	<b><i>Breed:</i></b>				
	Swedish Red and White	71	27	2	100
	Swedish Holstein	100	0	0	100
	Swedish Polled	25	0	0	25
20	Swedish Jersey	23	0	0	23

<sup>1</sup>Evaluation based on two trained persons. Class 2 = both person recognized a strong off-flavor (strong/strong); Class 1 = strong/moderate, moderate/moderate, moderate/normal.

- 25 ***The R238X substitution occurs at a surprisingly high frequency in the Swedish Red and White breed***  
 The population screening involved 248 animals representing the four dairy breeds in Sweden. The R238X substitution was not found in Swedish Holstein, Swedish Polled, or Swedish Jersey but was surprisingly common in the Swedish Red and White breed (Table 2). Two homozygous mutant and 27 heterozygotes were observed among 100 animals and the allele frequency estimate for the mutation was as high as  
 30 15.5%.

SNPs observed by cDNA sequencing of a heterozygous R238X carrier and a homozygous normal animal

- Only the wild type transcript appeared well represented in the mRNA sample from the R238X carrier. Only a very weak sequencing signal could be observed from the mutant transcript, at least 10 times weaker than  
 35 from the wild type transcript. A similar result was obtained using liver mRNA from a second heterozygous R238X carrier. Consequently, we could not search for possible additional mutations present in the *FM03* allele associated with fishy-off flavour. However, the RT-PCR experiment confirmed that the R238X mutation occurs in an expressed *FM03* gene.

- The comparison of the full-length liver cDNA sequence from the normal allele from the carrier and  
 40 from the homozygous normal individual revealed three additional single nucleotide polymorphisms (SNPs), all compiled in Table 3. Only one of the SNPs altered the amino acid sequence, E287G.

Table 3. Single nucleotide polymorphisms in the bovine *FMO3* gene

5	Nucleotide/Amino acid position <sup>1</sup>	Nucleotide/Amino acid change
	877/287	A↔G / E↔G
	1067/350	T↔C / no change
	1085/356	A↔C / no change

10

<sup>1</sup>Position as numbered in SEQ ID NO: 14.

#### Discussion

The present study has provided compelling evidence for that the observed *FMO3* nonsense mutation R238X causes fishy off-flavor in milk. This conclusion is based on (i) the identification of an obvious loss-of-function mutation in a gene associated with a similar syndrome in humans, (ii) a very strong association between the presence of this mutation and a strong fishy-off flavor in a case/control material and finally (iii) the presence of this mutation in the only breed (SRB) tested with well documented cases of a fishy off-flavor.

The R238X mutation causes a premature termination eliminating more than 50% of the *FMO3* enzyme which is expected to comprise 532 amino acids as in the human *FMO3* protein. Similar nonsense mutations in humans (E305X, E314X) have been shown to lead to a complete loss of enzyme activity (Treacy *et al.* 1998, Akerman *et al.* 1999). Moreover, we observed only barely detectable levels of the mutant transcript in liver mRNA from two heterozygous carriers of R238X. This is most probably explained by nonsense-mediated mRNA decay (NMD; Culbertson 1999). It is very likely that cows homozygous for the R238X mutation show a complete lack of *FMO3* activity necessary for the oxidation of TMA to an odorless compound. This provides a plausible explanation for the fishy off-flavor of the milk from these cows. Milk samples from five of the cows homozygous for the mutation and included in this study have previously been analyzed for TMA content using dynamic headspace gas chromatography. The results showed that these milk samples all had comparatively high concentrations of TMA whereas the milk from normal cows showed non-detectable or, in a few cases, very low TMA concentrations.

The results suggested a recessive mode of inheritance for the fishy-off flavor in cows milk in agreement with the recessive inheritance of the fishy-odor syndrome in humans. However, there was not a perfect agreement between this model and the observations from the case/control study since one carrier was scored as having strong fishy off-flavor (Class 2) and one homozygous mutant showed a moderate off-flavor (Class 1). We think there are two potential explanations from this deviation from a strict recessive model. The fishy-off flavor is influenced by environmental factors, like the presence of TMA precursors or *FMO* inhibitors in the feed. For example, the TMA concentration in the milk from one of the homozygous mutants was shown to vary between 1 and 16 mg TMA/kg milk within a single lactation. Secondly, the sensory evaluation is a somewhat subjective measure. The test panel consisting of two trained persons goes through a considerable number of milk samples during one day and may have some difficulties to discern the various off-flavors present in the milk samples and also to discriminate between different flavors.

In the present material the mutation was only observed among the SRB animals, but at a relatively high frequency. This is in complete accordance with the experiences made by the staff at the milk analysis laboratory that performed the sensory analysis, who have so far only encountered the fishy off-flavor in milk from SRB cows. We analyzed the pedigree data for all carriers and homozygous mutants but we did not find any common ancestor of these animals within the last 10 generations. The results suggest that the

mutation has been present in the existing SRB population for a relatively long period of time. The origin of the mutation probably dates back at least to the beginning of the last century, judging from the available pedigree information on affected animals. Moreover, the pedigree data indicate that the mutation may exist in other cattle populations closely related to the Ayrshire.

- 5 To the best of our knowledge, this is the first identified gene that has a profound influence on perceived quality of raw milk. The genotyping method described in this study can now be used by the breeding organisations to eliminate the problem in those breeds where a *FMO3* nonsense mutation is present. A practical way to reduce the problem would be to genotype currently used breeding bulls, bull dams, and young potential breeding bulls before progeny testing and eliminate carriers from breeding.

10

Example 3. Sequencing of the chicken *FMO3* gene

- Exonic parts of the chicken *FMO3* gene will be amplified from genomic DNA samples using both primers corresponding to sequences well conserved between the bovine, human, rabbit, mouse, and rat *FMO3* sequences as well as primers corresponding to EST sequences derived from the chicken *FMO3* gene, ESTs  
 15 603149708F1 and 603610105F1 available in the Chicken EST database ([http://www.chick.umist.ac.uk/cgi-bin/chicken\\_database.cgi](http://www.chick.umist.ac.uk/cgi-bin/chicken_database.cgi)). Both genomic DNA samples from hens that have been shown to produce eggs with a fishy off-flavor and genomic DNA samples from normal hens will be used as templates. The sequence of cDNA coding for the chicken *FMO3* will be obtained from RT-PCR products from liver mRNA.  
 The sequences obtained from hens that have been shown to produce eggs with a fishy off-flavor and  
 20 sequences obtained from normal hens will be compared and sequence variants identified.

Hens will be genotyped and *FMO3* variants associated with production of eggs with fishy off-flavour will be identified.

## REFERENCES

- Akerman, B. R., Forrest, S., Chow, L., Youil, R., Knight, M. & Treacy, E. P. 1999. Two novel mutations of the FMO3 gene in a proband with trimethylaminuria. *Human Mutation* 13:376-379
- 5 Al-Walz, M., Mitchell, S. C., Idle, J. R. & Smith, R. L. 1987a. The metabolism of <sup>14</sup>C-labelled trimethylamine and its N-oxide in man. *Xenobiotica* 17:551-558
- Al-Walz, M., Ayesh, R., Mitchell, S. C., Idle, J. R. & Smith, R. L. 1987b. Trimethylaminuria (fish-odour syndrome) - An inborn error of oxidative metabolism. *The Lancet* 8533:634-635
- Al-Walz, M., Ayesh, R., Mitchell, S. C., Idle, J. R. & Smith, R. L. 1987c. A genetic polymorphism of the N-oxidation of trimethylamine in humans. *Clinical Pharmacology and Therapeutics* 42:588-594
- 10 Al-Walz, M., Ayesh, R., Mitchell, S. C., Idle, J. R. & Smith, R. L. 1989. Trimethylaminuria: the detection of carriers using a trimethylamine load test. *Journal of Inherited Metabolic Disease* 12:80-85
- Ayesh, R. & Smith, R. L. 1990. Genetic polymorphism of trimethylamine oxidation. *Pharmacology & Therapeutics* 45:387-401
- 15 Ayesh, R., Mitchell, S. C., Zhang, A. & Smith, R. L. 1993. The fish odour syndrome: biological, familial, and clinical aspects. *British Medical Journal* 307: 655-657
- Basarab, T. B., Ashton, G. H. S., Menagé, H. duP. & McGrath, J. A. 1999. Sequence variation in the flavin-containing mono-oxygenase 3 gene (FMO3) in fish odour syndrome. *British Journal of Dermatology* 140: 164-167
- 20 Bolton, W., Carter, T. C. & Morley Jones, R. 1976. The hen's egg: Genetics of taints in eggs from hens fed on rapeseed meal. *British Poultry Science* 17: 313-320
- Corfield, A. 1955. Bacterial fishiness in milk. *Dairy Industries* December: 1035-1037
- Culbertson, M.R. 1999. RNA surveillance unforeseen consequences for gene expression, inherited genetic disorders and cancer. *Trends in Genetics* 15 :74-80
- 25 Dolphin, C. T., Riley, J. H., Smith, R. L., Shepard, E. A. & Phillips, I. R. 1997a. Structural organization of the human flavin-containing monooxygenase 3 gene (FMO3), the favoured candidate for fish-odor syndrome, determined directly from genomic DNA. *Genomics* 46: 260-267
- Dolphin, C. T., Janmohamed, A., Smith, R. L., Shepard, E. A. & Phillips, I. R. 1997b. Missense mutation in flavin-containing mono-oxygenase 3 gene, FMO3, underlies fish-odour syndrome. *Nature genetics* 17: 491-494
- 30 Eyer, H., Gauch, R. & Bosset, J. O. 1990. Untersuchung des Fehlgeruchs "fischig" in Kaffeerahm [The fishy off-flavour of coffee cream]. *Schweizerische Milchwirtschaftliche Forschung* 19 :43-45
- Forrest, S. M., Knight, M., Akerman, B. R., Cashman, J. R. & Treacy, E. P. 2001. A novel deletion in the flavin-containing monooxygenase gene (FMO3) in a Greek patient with trimethylaminuria. *Pharmacogenetics* 11: 169-174
- 35 Hlavica, P. & Kehl, M. 1977. Studies on the mechanism of hepatic microsomal N-oxide formation, the role of cytochrome P-450 and mixed function amine oxidase in the N-oxidation of N,N-dimethylamine. *Biochemical Journal* 164: 487-496
- Hobson-Frohock, A., Land, D. G., Griffiths, N. M. & Curtis, R. F. 1973. Egg taints: association with trimethylamine. *Nature* 243: 304-305
- 40 Humbert, J. R., Hammond, K. B., Hathaway, W. E., Marcoux, J. G. & O'Brien, D. 1970. Trimethylaminuria: the fish-odour syndrome. *The Lancet* 2: 770-771
- Humphriss, E. 1953. Chromogenic bacteria and milk. *Dairy Industries* September :781-782
- Kim, H. S., Gilliland, S. E., & Von Gunten, R. L. 1980. Chemical test for detecting wheat pasture flavor in cow's milk. *Journal of Dairy Science* 63:368-374
- 45 Metha, R. S., Bassette, R. & Ward, G. 1974. Trimethylamine responsible for fishy flavor in milk from cows on wheat pasture. *Journal of Dairy Science* 57:285-289
- Pearson, A. W., Butler, E. J., Curtis, R. F., Fenwick, G. R., Hobson-Frohock, A. & Land, D. G. 1979. Effect of rapeseed meal on trimethylamine metabolism in the domestic fowl in relation to egg taint. *Journal of the Science of Food and Agriculture* 30:799-804
- 50 Pearson, A. W., Greenwood, N. M., Butler, E. J. & Fenwick, G. R. 1982. The effects of thionamides and related compounds on trimethylamine oxidase activity in hepatic microsomes isolated from chickens (*Gallus domesticus*). *Comparative Biochemistry and Physiology. C: Comparative Pharmacology* 73C: 389-393

- Ronaghi, M., Uhlén, M. & Nyrén, P. 1998. A sequencing method based on real-time pyrophosphate. *Science* 281: 363, 365.
- Spellacy, E., Watts, R W. E. & Gollamall, S. K. 1979. Trimethylaminuria. *Journal of Inherited Metabolic Disease* 2: 85-88
- 5 Swoboda, P. A. T., & Peers, K. E. 1977. Volatile odorous compounds responsible for metallic, fishy taint formed in butterfat by selective oxidation. *Journal of the Science of Food and Agriculture* 28: 1010-1018
- Treacy, E. P., Akerman, B. R., Chow, L. M. L., Youll, R., Bibeau, C., Lin, J., Bruce, A. G., Knight, M., Danks, D. M., Cashman, J. R. & Forrest, S. M. 1998. Mutations of the flavin-containing monooxygenase gene (*FMO3*) cause trimethylaminuria, a defect in detoxication. *Human Molecular Genetics* 7: 839-845
- 10 Von Gunten, R. L., Bush, L. J., Odell, G. V., Wells, M. E. & Adams, G. D. 1976. Factors related to the occurrence of trimethylamine in milk. *Journal of Milk and Food Technology* 39 526-529
- Zhang, A.Q., Mitchell, S.C. & Smith, R.L. 1999. Dietary precursors of trimethylamine in man: a pilot study. *Food and Chemical Toxicology* 37: 515-520



## SEQUENCE LISTING

5 <110> Lunden, Anne  
Andersson, Leif  
Marklund, Stefan  
10  
  
<120> Mutations in the FMO3 gene  
15  
  
<130> Ref  
  
20  
<160> 20  
  
25 <170> PatentIn version 3.1  
  
30 <210> 1  
<211> 20  
<212> DNA  
35 <213> Artificial Sequence  
  
40 <220>  
<223> Primer  
<400> 1  
gaccatgcag aagagggcag  
45  
  
50 <210> 2  
<211> 27  
<212> DNA  
<213> Artificial Sequence  
55  
  
<220>  
<223> Primer

20

<400> 2  
 cttaaactgt atgtatttca ggaggtt 27  
 5  
 <210> 3  
 <211> 21  
 10 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Primer  
 20 <400> 3  
 catcatcagc tccagaagtg g 21  
 <210> 4  
 25 <211> 21  
 <212> DNA  
 30 <213> Artificial Sequence  
 <220>  
 35 <223> Primer  
 <400> 4  
 taaaggcatc aagccatagt t 21  
 40  
 <210> 5  
 <211> 21  
 45 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Primer  
 55 <400> 5  
 cagaatcctg aggaaagagc c 21

<210> 6  
 <211> 20  
 5 <212> DNA  
 <213> Artificial Sequence  
  
 10  
 <220>  
 <223> Primer  
 15 <400> 6  
 attacttggtg ctgcccagcg  
  
 20  
 <210> 7  
 <211> 26  
 <212> DNA  
 25 <213> Artificial Sequence  
  
 30  
 <220>  
 <223> Primer  
 <400> 7  
 gatgaatgat attaatgaga aaatgg  
 35  
 <210> 8  
 <211> 18  
 40 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> Primer  
 50 <400> 8  
 ccggtccac tgggtcag  
 55 <210> 9  
 <211> 142

20

26

18

<212> DNA

<213> Bos taurus

5

<220>

<221> exon

10

<222> (2)..(142)

<223>

15

<400> 9

20 g gcc agc att tat cgg tca gtc ttt acc aac tct tcc aaa gag atg aca 49  
Ala Ser Ile Tyr Arg Ser Val Phe Thr Asn Ser Ser Lys Glu Met Thr  
1 5 10 15

25 tgt ttt cca gac ttt cca ttt cct gat gat ttt cct aac ttt atg cac 97  
Cys Phe Pro Asp Phe Pro Phe Pro Asp Asp Phe Pro Asn Phe Met His  
20 25 30

30 aac agc aag ctc cag gaa tat att act atg ttt gcc aaa gaa aag 142  
Asn Ser Lys Leu Gln Glu Tyr Ile Thr Met Phe Ala Lys Glu Lys  
35 40 45

<210> 10

<211> 151

35

<212> DNA

<213> Bos taurus

40

<220>

<221> exon

45

<222> (2)..(151)

<223>

50

<400> 10

55 c tcc tgg gtg atg agc cgg gtc tgg gat gaa ggc tat cca tgg gac atg 49  
Ser Trp Val Met Ser Arg Val Trp Asp Glu Gly Tyr Pro Trp Asp Met  
1 5 10 15

ctg ttt atc act cga ttt gaa aca ttc ctc aag aac acc tta ccg aca 97  
Leu Phe Ile Thr Arg Phe Glu Thr Phe Leu Lys Asn Thr Leu Pro Thr  
20 25 30

gtc att tct aac tgg tgg tac atg aag caa atg aac gcc aga ttc aag 145  
 Val Ile Ser Asn Trp Trp Tyr Met Lys Gln Met Asn Ala Arg Phe Lys  
 35 40 45  
 5  
 cac gag 151  
 His Glu  
 50  
 10  
 <210> 11  
 <211> 313  
 15 <212> DNA  
 <213> Bos taurus  
 20  
 <220>  
 <221> exon  
 25 <222> (2)..(313)  
 <223>  
 30  
 <400> 11  
 t gtg ttc aat gac gaa ctc cca gct tgc att tta tgt ggc att gtg acc 49  
 Val Phe Asn Asp Glu Leu Pro Ala Cys Ile Leu Cys Gly Ile Val Thr  
 1 5 10 15  
 35  
 att aag cca aat gtg aag gag ttt aca gag gat tca gct att ttt gag 97  
 Ile Lys Pro Asn Val Lys Glu Phe Thr Glu Asp Ser Ala Ile Phe Glu  
 20 25 30  
 40  
 gat ggg acg gtg ttt aag gcc att gac tat gtc atc ttt gca aca ggc 145  
 Asp Gly Thr Val Phe Lys Ala Ile Asp Tyr Val Ile Phe Ala Thr Gly  
 35 40 45  
 tat agt tat gcc tac ccc ttc ctt gat gac tcc atc att aag agc aga 193  
 45 Tyr Ser Tyr Ala Tyr Pro Phe Leu Asp Asp Ser Ile Ile Lys Ser Arg  
 50 55 60  
 gac aat gag gtc acc tta ttt aaa ggc ata ttc cca cct cca ctg gaa 241  
 Asp Asn Glu Val Thr Leu Phe Lys Gly Ile Phe Pro Pro Pro Leu Glu  
 50 65 70 75 80  
 aag cca acc ttg gct gtg atc ggc ctt gtc cag tcc ctt gga gct gcc 289  
 Lys Pro Thr Leu Ala Val Ile Gly Leu Val Gln Ser Leu Gly Ala Ala  
 85 90 95  
 55  
 atc ccc act act gac ctg cag tct 313  
 Ile Pro Thr Thr Asp Leu Gln Ser  
 100

<210> 12  
 <211> 202  
 5 <212> DNA  
 <213> Bos taurus  
 10  
 <220>  
 <221> exon  
 15 <222> (2)..(202)  
 <223>  
 20  
 <400> 12  
 g ttt ggc aaa agt gat acc ata cag acg gat tat gtt gtt tat atg gat 49  
 Phe Gly Lys Ser Asp Thr Ile Gln Thr Asp Tyr Val Val Tyr Met Asp  
 25 1 5 10 15  
 gag ctt gcc tcc ttc att ggg gca aag ccc aac atc cca tgg ctg ttt 97  
 Glu Leu Ala Ser Phe Ile Gly Ala Lys Pro Asn Ile Pro Trp Leu Phe  
 20 25 30  
 30 ctc aca gat cca aag ttg gca ttg gag gtc tac ttt ggc cct tgc acc 145  
 Leu Thr Asp Pro Lys Leu Ala Leu Glu Val Tyr Phe Gly Pro Cys Thr  
 35 40 45  
 35 cca tac cag ttt agg ctg gtg ggc cca gga aag tgg cca gga gcc aga 193  
 Pro Tyr Gln Phe Arg Leu Val Gly Pro Gly Lys Trp Pro Gly Ala Arg  
 50 55 60  
 aac gcc atc  
 40 Asn Ala Ile 202  
 65  
 <210> 13  
 45 <211> 714  
 <212> DNA  
 50 <213> Bos taurus  
 <220>  
 55 <221> Intron  
 <222> (1)..(714)

<223>

5 <220>

<221> Intron

<222> (1)..(714)

10

<223>

15 <400> 13

ctcagattgt aagagcatct gtggtaaagg ggatccagaa ttttcatagc aaaatttgac 60

tttgtgaata cttaaactgt tttagtcttg aatactcctg agtgggacca ttaggggtgac 120

20 agctatgagc taaattcagc cattcaacaa atagttatca aggatatact tcgcacctga 180

tagaccacta aatcctggtt cagccctcca tatctcctgc aatatatcat aaaagagaga 240

25 cccagaaaga ttaaatttca gaggagcttg ggaacacacg gatgtggact tctattttct 300

agctcataca tcctctaaaa tgtaaggatg ccttccaata ggtggctcta ataaatgact 360

cagccacaca acaactagtc agaagaacta ggctccagtc ctacctctgt cactaatcag 420

30 ctcagtggcc atgaattaat aactacacgt gagcaagaga caattcctct acttgtaaatt 480

gtggataaaa atgcctttct cagcagcctt gttgtgttgt ggaatgtcca agtgagatca 540

35 gtatgaaagt ccttcagtaa aatctacaaa gtactggata atgagcaata atccccttac 600

caacaaaagg atcgatgttg atttgcgtgg ataaaaagggt gtgaggcatt ttccctgttc 660

ttgtttctaa aggaagctcg gatagccaca gtcttgtttc tctccctcct ctag 714

40

<210> 14

<211> 1672

45 <212> DNA

<213> Bos taurus

50

<220>

<221> CDS

55 <222> (18)..(1616)

<223>

&lt;220&gt;

5 &lt;221&gt; mRNA

&lt;222&gt; (1)..(1672)

&lt;223&gt;

10

&lt;400&gt; 14

15	attacagaaa ggtaact atg gtg aag aaa gtg gcc atc att gga gca ggc	50
	Met Val Lys Lys Val Ala Ile Ile Gly Ala Gly	
	1 5 10	
20	atc agt ggc ctg gcc tcc atc aga aac tgc cta gaa gag gga ctg gaa	98
	Ile Ser Gly Leu Ala Ser Ile Arg Asn Cys Leu Glu Glu Gly Leu Glu	
	15 20 25	
25	ccc acc tgc ttt gag aag ggt gaa gac att ggg ggc ctg tgg aaa ttc	146
	Pro Thr Cys Phe Glu Lys Gly Glu Asp Ile Gly Gly Leu Trp Lys Phe	
	30 35 40	
30	tgc gac cat gta gag gaa ggc agg gcc agc att tat cgg tca gtc ttt	194
	Ser Asp His Val Glu Glu Gly Arg Ala Ser Ile Tyr Arg Ser Val Phe	
	45 50 55	
35	acc aac tct tcc aaa gag atg aca tgt ttt cca gac ttt cca ttt cct	242
	Thr Asn Ser Ser Lys Glu Met Thr Cys Phe Pro Asp Phe Pro Phe Pro	
	60 65 70 75	
40	gat gat ttt cct aac ttt atg cac aac agc aag ctc cag gaa tat att	290
	Asp Asp Phe Pro Asn Phe Met His Asn Ser Lys Leu Gln Glu Tyr Ile	
	80 85 90	
45	act atg ttt gcc aaa gaa aag aac ctc ctg aaa tac ata caa ttt aag	338
	Thr Met Phe Ala Lys Glu Lys Asn Leu Leu Lys Tyr Ile Gln Phe Lys	
	95 100 105	
50	aca att gta tcc agt gta aat aag cgt ccc gat ttc caa acc act ggc	386
	Thr Ile Val Ser Ser Val Asn Lys Arg Pro Asp Phe Gln Thr Thr Gly	
	110 115 120	
55	caa tgg gat gtt atc act gaa aag gat ggt aaa aag gaa tca gct gtc	434
	Gln Trp Asp Val Ile Thr Glu Lys Asp Gly Lys Lys Glu Ser Ala Val	
	125 130 135	
60	ttt gat gcc gta atg att tgt tct gga cat cat gtg tac ccc aac ata	482
	Phe Asp Ala Val Met Ile Cys Ser Gly His His Val Tyr Pro Asn Ile	
	140 145 150 155	
65	cct aaa gag tcc ttt cca gga ata aaa ctt ttt aaa ggc aaa tgc ttc	530
	Pro Lys Glu Ser Phe Pro Gly Ile Lys Leu Phe Lys Gly Lys Cys Phe	
	160 165 170	
70	cac agc cgg gac tat aaa gaa cca gga atc ttc aag ggg aag cga gtc	578
	His Ser Arg Asp Tyr Lys Glu Pro Gly Ile Phe Lys Gly Lys Arg Val	



	175	180	185	
5	ctg gtg att ggt ctg ggg aac tca ggc tgt gac atc gcc tca gaa ctc Leu Val Ile Gly Leu Gly Asn Ser Gly Cys Asp Ile Ala Ser Glu Leu 190 195 200			626
10	agc cac ata gct gaa aaa gtc atc atc agc tcc cga agt ggc tcc tgg Ser His Ile Ala Glu Lys Val Ile Ile Ser Ser Arg Ser Gly Ser Trp 205 210 215			674
	gtg atg agc cgg gtc tgg gat gaa ggc tat cca tgg gac atg ctg ttt Val Met Ser Arg Val Trp Asp Glu Gly Tyr Pro Trp Asp Met Leu Phe 220 225 230 235			722
15	atc act cga ttt gaa aca ttc ctc aag aac acc tta ccg aca gtc att Ile Thr Arg Phe Glu Thr Phe Leu Lys Asn Thr Leu Pro Thr Val Ile 240 245 250			770
20	tct aac tgg tgg tac atg aag caa atg aac gcc aga ttc aag cac gag Ser Asn Trp Trp Tyr Met Lys Gln Met Asn Ala Arg Phe Lys His Glu 255 260 265			818
25	aac tac ggc ttg atg cct tta aac agc acc ctg agg aaa gag cct gtg Asn Tyr Gly Leu Met Pro Leu Asn Ser Thr Leu Arg Lys Glu Pro Val 270 275 280			866
30	ttc aat gac gaa ctc cca gct tgc att tta tgt ggc att gtg acc att Phe Asn Asp Glu Leu Pro Ala Cys Ile Leu Cys Gly Ile Val Thr Ile 285 290 295			914
	aag cca aat gtg aag gag ttt aca gag gat tca gct att ttt gag gat Lys Pro Asn Val Lys Glu Phe Thr Glu Asp Ser Ala Ile Phe Glu Asp 300 305 310 315			962
35	ggg acg gtg ttt aag gcc att gac tat gtc atc ttt gca aca ggc tat Gly Thr Val Phe Lys Ala Ile Asp Tyr Val Ile Phe Ala Thr Gly Tyr 320 325 330			1010
40	agt tat gcc tac ccc ttc ctt gat gac tcc atc att aag agc aga gac Ser Tyr Ala Tyr Pro Phe Leu Asp Asp Ser Ile Ile Lys Ser Arg Asp 335 340 345			1058
45	aat gag gtc acc tta ttt aaa ggc atc ttc cca cct cca ctg gaa aag Asn Glu Val Thr Leu Phe Lys Gly Ile Phe Pro Pro Pro Leu Glu Lys 350 355 360			1106
50	cca acc ttg gct gtg atc ggc ctt gtc cag tcc ctt gga gct gcc atc Pro Thr Leu Ala Val Ile Gly Leu Val Gln Ser Leu Gly Ala Ala Ile 365 370 375			1154
	ccc act act gac ctg cag tct cgc tgg gca gta caa gta att aag gga Pro Thr Thr Asp Leu Gln Ser Arg Trp Ala Val Gln Val Ile Lys Gly 380 385 390 395			1202
55	aca tgc cct ttg cct tct gtc aag gac atg atg aat gat att gat gaa Thr Cys Pro Leu Pro Ser Val Lys Asp Met Met Asn Asp Ile Asp Glu 400 405 410			1250
	aaa atg ggg aaa aag ctc aaa ttg ttt ggc aaa agt gat acc ata cag			1298

Lys Met Gly Lys Lys Leu Lys Leu Phe Gly Lys Ser Asp Thr Ile Gln  
 415 420 425  
 5 acg gat tat gtt gtt tat atg gat gag ctt gcc tcc ttc att ggg gca 1346  
 Thr Asp Tyr Val Val Tyr Met Asp Glu Leu Ala Ser Phe Ile Gly Ala  
 430 435 440  
 10 aag ccc aac atc cca tgg ctg ttt ctc aca gat cca aag ttg gca ttg 1394  
 Lys Pro Asn Ile Pro Trp Leu Phe Leu Thr Asp Pro Lys Leu Ala Leu  
 445 450 455  
 gag gtc tac ttt ggc cct tgc acc cca tac cag ttt agg ctg gtg ggc 1442  
 Glu Val Tyr Phe Gly Pro Cys Thr Pro Tyr Gln Phe Arg Leu Val Gly  
 460 465 470 475  
 15 cca gga aag tgg cca gga gcc aga aac gcc atc ctg acc cag tgg gac 1490  
 Pro Gly Lys Trp Pro Gly Ala Arg Asn Ala Ile Leu Thr Gln Trp Asp  
 480 485 490  
 20 cgg tta ctg aaa cct atg acg aca aga gtg gtt ggg agt cct ctg aag 1538  
 Arg Leu Leu Lys Pro Met Thr Thr Arg Val Val Gly Ser Pro Leu Lys  
 495 500 505  
 25 cct tgc tta ttt tgc aac tgg ttc aga cct gtt ctt att tct gtt gta 1586  
 Pro Cys Leu Phe Cys Asn Trp Phe Arg Pro Val Leu Ile Ser Val Val  
 510 515 520  
 30 tca att gct gct ctc att gtg ttg ttc tag ccatcattct atctaggatt 1636  
 Ser Ile Ala Ala Leu Ile Val Leu Phe  
 525 530  
 ctgaacatta ctaacaatac ctggaaagaa gcttca 1672  
 35 <210> 15  
 <211> 532  
 <212> PRT  
 40 <213> Bos taurus  
 45 <400> 15  
 Met Val Lys Lys Val Ala Ile Ile Gly Ala Gly Ile Ser Gly Leu Ala  
 1 5 10 15  
 50 Ser Ile Arg Asn Cys Leu Glu Glu Gly Leu Glu Pro Thr Cys Phe Glu  
 20 25 30  
 55 Lys Gly Glu Asp Ile Gly Gly Leu Trp Lys Phe Ser Asp His Val Glu  
 35 40 45  
 Glu Gly Arg Ala Ser Ile Tyr Arg Ser Val Phe Thr Asn Ser Ser Lys

Condition	Group 55 (%)	Group 25 (%)	Group 22 (%)
1	85	75	65
2	75	65	55
3	70	60	50
4	75	65	55
5	65	70	60

55



5

10

15

15

20

25

30

35

35

40

45

50

55

55

Pro Ala Cys Ile Leu Cys Gly Ile Val Thr Ile Lys Pro Asn Val Lys  
 290 295 300

5 Glu Phe Thr Glu Asp Ser Ala Ile Phe Glu Asp Gly Thr Val Phe Lys  
 305 310 315 320

10 Ala Ile Asp Tyr Val Ile Phe Ala Thr Gly Tyr Ser Tyr Ala Tyr Pro  
 325 330 335

15 Phe Leu Asp Asp Ser Ile Ile Lys Ser Arg Asp Asn Glu Val Thr Leu  
 340 345 350

20 Phe Lys Gly Ile Phe Pro Pro Pro Leu Glu Lys Pro Thr Leu Ala Val  
 355 360 365

25 Ile Gly Leu Val Gln Ser Leu Gly Ala Ala Ile Pro Thr Thr Asp Leu  
 370 375 380

30 Gln Ser Arg Trp Ala Val Gln Val Ile Lys Gly Thr Cys Pro Leu Pro  
 385 390 395 400

35 Ser Val Lys Asp Met Met Asn Asp Ile Asp Glu Lys Met Gly Lys Lys  
 405 410 415

40 Leu Lys Leu Phe Gly Lys Ser Asp Thr Ile Gln Thr Asp Tyr Val Val  
 420 425 430

45 Tyr Met Asp Glu Leu Ala Ser Phe Ile Gly Ala Lys Pro Asn Ile Pro  
 435 440 445

50 Trp Leu Phe Leu Thr Asp Pro Lys Leu Ala Leu Glu Val Tyr Phe Gly  
 450 455 460

55 Pro Cys Thr Pro Tyr Gln Phe Arg Leu Val Gly Pro Gly Lys Trp Pro  
 465 470 475 480

60 Gly Ala Arg Asn Ala Ile Leu Thr Gln Trp Asp Arg Leu Leu Lys Pro  
 485 490 495

65 Met Thr Thr Arg Val Val Gly Ser Pro Leu Lys Pro Cys Leu Phe Cys  
 500 505 510

70 Asn Trp Phe Arg Pro Val Leu Ile Ser Val Val Ser Ile Ala Ala Leu  
 515 520 525

Ile Val Leu Phe  
530

5  
 <210> 16  
 <211> 21  
 10 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> Primer  
 20 <400> 16  
 gatgaaggct atccatggga c  
 25 <210> 17  
 <211> 24  
 <212> DNA  
 30 <213> Artificial Sequence  
 35 <220>  
 <223> Primer  
 <400> 17  
 40 taaaggcatc aagccgtagt tctc  
 <210> 18  
 45 <211> 18  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 55 <223> Primer  
 <400> 18  
 tgaggaatgt ttcaaadc

21

24

18

<210> 19  
<211> 27  
5 <212> DNA  
<213> Artificial Sequence  
10  
<220>  
<223> Primer  
15 <400> 19  
ggacttagac acacagaaga aaagaag  
20 <210> 20  
<211> 29  
<212> DNA  
25 <213> Artificial Sequence  
  
30 <220>  
<223> Primer  
<400> 20  
35 gaggtgtgaa attcttattt tttaaataag

27

29

## CLAIMS

1. A polypeptide which is a flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide comprising at least a sequence having at least 85% identity with the polypeptide sequence SEQ ID NO: 15.
- 5 2. A polypeptide which is a functionally altered mutant of flavin-containing monooxygenase 3 (FMO3), wherein said FMO3 is a polypeptide comprising at least a sequence having at least 85% identity with the polypeptide sequence SEQ ID NO: 15.
- 10 3. A polypeptide according to claim 2 resulting from a deletion, insertion, non-sense, or mis-sense mutation in a FMO3 gene..
4. A polypeptide according to claim 3 which is the R238X variant of the bovine FMO3.
- 15 5. An isolated nucleic acid sequence encoding a polypeptide according to any of claims 1-4, or the complement thereof.
6. An isolated nucleic acid sequence according to claim 5 which is the nucleic acid sequence shown in SEQ ID NO: 14.
- 20 7. A nucleic acid sequence comprising at least a portion of a nucleic acid sequence encoding a polypeptide of any of claims 1-4, and up to 500 kb of a 3' and/or a 5' adjacent genomic DNA sequence, or the complement thereof.
- 25 8. A nucleic acid fragment selected from the group consisting of;
- a specific fragment of a nucleic acid sequence encoding a polypeptide according to any claims 1-4,
  - a nucleic which specifically hybridises under stringent condition with a nucleic acid sequence encoding a polypeptide according to any of claims 1-4,
  - a specific fragment of a nucleic acid sequence according to claim 7,
  - 30 - a nucleic which specifically hybridises under stringent condition with a nucleic acid sequence according to claim 7,
  - SEQ ID NO: 9,
  - SEQ ID NO: 10,
  - SEQ ID NO: 11,
  - 35 - SEQ ID NO: 12,
  - SEQ ID NO: 16,
  - SEQ ID NO: 17, and
  - SEQ ID NO: 18.
- 40 9. A set of primers for amplifying a nucleic acid sequence according to any of claims 5-7, comprising at least one primer consisting of a nucleic acid fragment according to claim 8.
10. A method for detecting a mutation in the *FMO3* gene of an animal where the mutation will cause an alteration in the metabolism of trimethylamine leading to a fish off-flavour in a food product of the animal
- 45 or its offspring, the wherein the method comprises:

- obtaining a sample of genomic DNA from the animal,
- amplifying a segment of said DNA spanning a polymorphic marker by PCR using a set of primers according to claim 9, and
- detecting in said amplified DNA the presence of an allele of a polymorphic marker associated with said mutation in the *FMO3* gene.

11. A method for detecting a mutation in the *FMO3* gene of an animal where the mutation will cause an alteration in the metabolism of trimethylamine leading to a fish off-flavour in a food product of the animal or its offspring, wherein the method comprises:

- obtaining a nucleic acid sample from the animal;
- determining the presence in said nucleic acid sample of a nucleic acid sequence encoding a mutated *FMO3*.

12. A method for detecting a nucleic acid sequence comprising a mutation in the *FMO3* gene of an animal where the mutation will cause an alteration in the metabolism of trimethylamine leading to a fish off-flavour in a food product of the animal or its offspring, wherein the method comprises:

- obtaining a nucleic acid sample from the animal;
- contacting said nucleic acid sample with a nucleic acid probe spanning said mutation under conditions of specific hybridisation between said probe and the mutant sequence to be detected
- detecting the hybridisation complex.

13. A method according to any of claims 11-12 wherein the presence of the nucleic acid sequence encoding said mutant polypeptide is determined by contacting the nucleic acid sample with a nucleic acid fragment according to claim 8.

14. A method according to any of claims 11-12 which further comprises PCR amplification from the nucleic acid sample, of a sequence comprising at least the portion of the *FMO3* sequence wherein the mutation is to be detected.




# Figure 1

Modtaget PVS

- 9 JAN. 2002

a)                    IleThrArg  
R238 : 5' ATCACTCGA  
R238X: 5' ATCACTTGA  
                      IleThrSTP

b)



BEST AVAILABLE COPY